

From the Association of Diabetes Care & Education Specialists

An Effective Model of Diabetes Care and Education: The ADCES7 Self-Care Behaviors™

Position Statement

Association of Diabetes Care and Education Specialists

From the Association of Diabetes Care and Education Specialists, Chicago, Illinois.

Correspondence to Leslie Kolb, Chief Science and Practice Officer, Association of Diabetes Care & Education Specialists, 125 S Wacker Dr, Suite 600, Chicago, IL 60606, USA (lkolb@adces.org).

Acknowledgments: ADCES gratefully acknowledges the following individuals for their role in creating this document:

Carole Mensing, RN, MA, CDCES (Co-Chair)

Donna Tomky, MSN, RN, C-NP, CDCES (Co-Chair)

Lois Moss-Barnwell, MS, RDN, LDN, CDCES

Sheri R. Colberg, PhD

Patricia Davidson, DCN, RDN, LDN, CDCES

Kim Coy DeCoste, RN, MSN, MLDE, CDCES

Nancy D'Hondt, RPh, CDCES

Paulina N. Duker, MPH, RN, CDCES

Leslie Kolb, RN, MBA

Eliot LeBow, LCSW, CDCES

Melinda Maryniuk, RDN, LD, CDCES

Molly McElwee-Malloy, RN, CDCES

David K. Miller, RN, MSEd, CDCES

Malinda Peebles, RN, MS, CDCES

Joanne Rinker, MS, RDN, LDN, CDCES

Cecilia Sauter MS, RDN, CDCES

Michael B. See, MS, RCEP, CDCES

Evan Sisson, MHA, PharmD, CDCES

ADCES gratefully acknowledges the following individuals for their role in reviewing this document:

Martha Funnell, RN, MS, CDCES

Barb Schreiner, RN, BC-ADM, PhD, CDCES

Jane Dickinson, RN, CDCES, PhD

Healthy Eating: Hope Warshaw, MMSc, RDN, CDCES, BC-ADM

Being Active: Gary Scheiner, MS, CDCES

Monitoring: Janice MacLeod, MA, RDN, LD, CDCES

Taking Medications: Geoff Twigg, PharmD, CDCES, BCACP

Problem Solving: Leigh Gemmell, PhD

Reducing Risk: Lucille Hughes, DNP, RN, CDCES, BC-ADM

Healthy Coping: Mary de Groot, PhD, HSPP

ADCES Team Members: Matthew Eaton and Kirsten Yehl, MS, MLIS

Medical Writer: Lauren Bronich-Hall, MS, RDN, LDN, CDCES (Health Illuminations, LLC)

Author's Note: This is an official statement of the Association of Diabetes Care & Education Specialists (ADCES). ADCES is an interprofessional membership organization dedicated to improving prediabetes, diabetes and cardiometabolic care through innovative education, management and support.

DOI: 10.1177/0145721720978154

© 2021 The Author(s)

Purpose

The ADCES7 Self-Care Behaviors™ (ADCES7) is a robust framework for self-management of diabetes and other related conditions, such as prediabetes and cardio-metabolic diseases. It is the position of the Association of Diabetes Care and Education Specialists (ADCES) that at the cornerstone of diabetes self-management education and support, the ADCES7 is the framework for achieving behavior change that leads to effective self-management through improved behavior and clinical outcome measures. The ADCES7 model guides the health care team in effective person-centered collaboration and goal setting to achieve health-related outcomes and improved quality of life. Continued research and evidence are critical to expand this model and broaden its application to other chronic conditions. Given the advances in the science of diabetes management as well as diabetes self-management education and support, ADCES has evaluated the ADCES7 within the framework of these advances, including the digital and dynamic health care landscape.

Conclusion

This revised position statement blends the updates in research and ADCES's vision and expansion beyond diabetes to refresh the ADCES7 framework. This revision reflects the perspectives of all members of the health care team as they problem solve with individuals who are at risk for or who have diabetes and related conditions to achieve healthier outcomes.

Introduction

In early 2020, the American Association of Diabetes Educators (AADE) changed its name to the Association of Diabetes Care & Education Specialists (ADCES). This was a multiyear project to position diabetes educators with a broader role as integrators for clinical management, education, prevention, and support. The decision to retitle was made through an extensive research process by ADCES while working with a professional research and branding firm. The research included both qualitative and quantitative surveys of health care professionals, including those who work in the field of diabetes, ADCES members, and nonmembers. This process revealed that

the word *specialist* conveyed knowledge and expertise, both traits that support the expanded position of the ADCES beyond educator. To reflect the high level of expertise, the term *diabetes educator* was replaced with *diabetes care and education specialist* to recognize the role of the diabetes care and education specialist as a part of the care team. It followed that the AADE7 Self-Care Behaviors® should be renamed the ADCES7 Self-Care Behaviors™ to support a consistent message.

The Association for Diabetes Care & Education's ADCES7 framework provides an evidenced-based model for assessment, intervention, and evaluation of individuals and populations living with diabetes and other cardio-metabolic conditions.¹ Using the ADCES7 framework, diabetes care and education specialists partner with people living with diabetes and related conditions to support informed decision-making. Diabetes care and education specialists embrace a person-centered philosophy, incorporating a strengths-based approach and acknowledging the whole person in the context of the person's life and relationships. The diabetes care and education specialist focus includes not only diabetes care, education, and ongoing support of self-management but also related conditions such as obesity, prediabetes, diabetes-related complications, and cardiometabolic disease. Given the advances in the science of diabetes management as well as diabetes self-management education and support, ADCES has evaluated the ADCES7 within the framework of these advances, including the digital and dynamic health care landscape. This revised position statement blends research updates and ADCES's vision and expansion beyond diabetes to revise the ADCES7 framework and encompasses the varied perspective of the health care team.

ADCES7 Revision

Aligned with ADCES's vision of "optimal health and quality of life for persons with, affected by, or at risk for diabetes and chronic conditions,"² the ADCES7 Self-Care Behaviors are as follows:

- Healthy Coping
- Healthy Eating
- Being Active
- Taking Medication
- Monitoring
- Reducing Risk
- Problem Solving.

Background

To conduct this “ADCES7 revision,” ADCES assembled a task force to review the literature and previous documents³⁻⁹ in an effort to update the ADCES7 framework while preserving its original intent.

AADE was challenged by the Centers for Medicare & Medicaid Services (CMS) in 1997 to identify the unique outcome measures of diabetes self-management education.¹⁰ AADE convened a task force to determine what to measure, when to monitor, and how to manage chronic disease over its continuum as it related to diabetes education and care.¹¹ This task force defined the unique outcomes of diabetes education as “behavior change”¹¹ and identified 7 self-care behaviors that promote successful and effective diabetes self-management, known as the AADE7. The work of the original taskforce included mapping the original 15 content areas of the 1995 National Standards for Diabetes Self-Management Education (NSDSME),¹⁰ developing and testing tools to capture outcomes,¹² and reaching a consensus on Diabetes Self-Management Education (DSME) outcome measures using the AADE7 Self-Care Behaviors.⁵ This framework shifted the focus away from educational content delivery to an outcome-driven practice using person-centered and self-determined goals.^{9,13} Effective diabetes education needed to go beyond knowledge transfer; it needed to address and support behavior change and affect clinical and health-related outcomes.

The NSDSME continues to include the AADE7 framework in its updates,¹⁴⁻¹⁶ and the AADE7 has provided standardized nomenclature for assessment, identification of self-care-related problems and barriers, goal setting, problem-solving, documentation, measurement, evaluation, quality improvement, and policy making.⁹ Based on additional research and practice, diabetes self-management education and support has evolved beyond knowledge and behavior change to focus on quality of life and person-centered approaches to management, education, and care. These gains in knowledge, along with innovative technologies, have changed some of the demands of self-care.

Role of Technology

The use of technology has transformed the approach to diabetes self-care and implementation of the ADCES7 framework. Technology developed to support self-care includes medical devices such as glucose meters, insulin

pumps, continuous glucose monitors; digital therapeutics such as mobile apps, text messaging, electronic communications, and video conference platforms; and wearable technologies such as Fitbits and Apple watches. People with diabetes and other related conditions can receive health care services virtually, outside of the clinic/office, with the capabilities of the Internet and mobile devices. This alleviates barriers such as transportation or cost of travel. Technology has the power to synthesize information into a digestible format, resulting in simplified interpretation and application to self-management. When designed with the user in mind, these tools can also serve to engage, encourage, and motivate self-care. Diabetes care and education specialists help with technology selection, device training, data downloads, data evaluation, troubleshooting, and back-up plans for times when technology fails.¹⁷ By collaborating with diabetes care and education specialists, people with diabetes and related conditions can learn how to use these technological tools effectively, when available, to improve their clinical and quality-of-life outcomes.

Although long-term studies are needed to evaluate sustainability,¹⁸ technology-enabled health care delivery can help people with diabetes and related conditions optimize their outcomes.¹⁹ Diabetes care and education specialists can provide the element of human touch to identify appropriate candidates and tools, provide training, and facilitate ongoing use of these tools and the information they offer. Research shows the use of electronic health records and wearables, along with the resulting patient-generated health data, can improve clinical outcomes and engagement.²⁰⁻²³ Mobile health interventions for obesity and diabetes have promoted behavior change,²⁴ and technology-enabled diabetes support has demonstrated clinically significant results in clinical settings.^{4,25} In people with type 1 diabetes, studies have validated that the use of continuous glucose monitoring can increase time in range and lower risk for severe hypoglycemia.²⁶

Accompanying these benefits, technology also brings new challenges to the health care team. To stay current with the accelerated growth of technology, the members of the health care team must familiarize themselves with the technologies, learn the intricacies of new devices, and overcome their technology phobias. People living with or affected by diabetes and related conditions, along with their diabetes care and education specialists, must be able to prioritize data surplus from technologies to minimize time and resource burdens.




Figure 1. Transformation of the ADCES7 image.

Reproduced with permission of the Association of Diabetes Care & Education Specialists, 2020

New ADCES7 image. Although originally presented in a list format, the ADCES7 Self-Care Behaviors overlap in nature, specifically the knowledge and skills to master them, barriers associated with their mastery, and associated outcome measures. Accordingly, ADCES has revised the image associated with the ADCES7 to underscore the interrelatedness of these behaviors (Figure 1).

Advances in the science of DSMES emphasize the independent effect of the emotional burden of diabetes on metabolic and quality-of-life outcomes. Because Healthy Coping must begin before learning can occur, this behavior is centrally located at the core of the new image (Figure 1) to symbolize its significance in sustainable diabetes self-management. The inner ring contains Healthy Eating, Being Active, and Taking Medication. These behaviors often serve as the basis for care plans because they comprise what individuals with diabetes and related conditions undertake regularly as they self-manage their condition. The next ring, Monitoring, encircles these 4 self-care behaviors. By collecting personalized data, Monitoring helps convert some of the intangible components of diabetes into perceptible ones. The knowledge gained and the ability to use the information from Monitoring can drive behavior change. Equally important, the outer ring contains the less tangible self-care behaviors of Reducing Risk and Problem Solving, which greatly influence motivation, goal setting, and the

ability to transform goals into action. The updated image of the ADCES7 depicts this interconnectivity.

Examination and Validation of the ADCES7

Healthy Coping

Healthy coping, defined as “a positive attitude toward diabetes and self-management, positive relationships with others, and quality of life,”²⁷ is critical for mastery of the other 6 behaviors. Psychosocial factors that interfere with a person’s ability to self-manage the disease and achieve desired metabolic outcomes greatly influence diabetes and other related conditions.²⁸⁻³² Person-centered care contributes to positive health outcomes and psychological well-being.³³ Conversely, diabetes-related distress negatively affects the physical and emotional well-being of the person living with diabetes.³⁴ Diabetes-related distress, described as the emotional burden of diabetes, constant demands from diabetes self-management, possibility of developing complications, and lack of support and access to care,^{29,34-36} hinders self-care. People living with diabetes are also more prone to depression and anxiety^{32,37} as well as disordered eating and cognitive impairment.³⁷ These psychosocial factors reduce the ability to self-manage. An evaluation by a diabetes care and education specialist and other members

of the health care team and appropriate referrals to behavioral specialists are necessary to support people living with diabetes and related conditions. Ongoing evaluation and support are key components to making sustainable behavior changes, as reinforced by the term change from *diabetes self-management education* to *diabetes self-management education and support*.

Behaviors That Contribute to Healthier Outcomes

Increase Self-Efficacy

Self-efficacy, described as an individual's belief in his or her own ability,³⁸ is critical to self-care. Both depression and diabetes-related distress influence self-efficacy in people living with diabetes.^{39,40} Research has shown that higher levels of positive emotions, self-efficacy, and increased social support and attitudes toward self-care behaviors are associated with improved self-care in diabetes and cardiovascular disease.^{32,41-44} Indicators of psychological well-being, including optimism, positive affect, self-efficacy, and gratitude, are associated with better health outcomes in many chronic diseases.³⁵ People with mild to moderate symptoms of diabetes-related distress can benefit from a referral to DSMES.⁴⁵ Those with more severe distress should be referred to mental health professionals prior to diabetes education.

Address Cognitive Impairment

Any impairment to learning, memory, attention, mental flexibility, and executive function can decrease the ability to perform self-care behaviors, resulting in inconsistent diabetes self-management and associated glyce-mic outcomes.^{30,46-50} Cognitive impairment is associated with type 1 and type 2 diabetes.⁵⁰ Mental health issues, such as diabetes distress, attention-deficit/hyperactivity disorder, depression, and addiction, also cause cognitive impairment. Cognitive impairment affects knowledge and skill transfer and the ability to learn and apply new information (Problem Solving).⁵¹

Gather Support

Diabetes care and education specialists, trained community health workers, family, and friends can provide support to people living with diabetes and related conditions. Together, people with diabetes and related conditions and their team members can identify effective

support networks to assist with changes in diabetes and life that occur over time. This support can also be virtual; technology has introduced options such as online peer support and telehealth support to help people share their concerns and feelings around diabetes self-management. Peer support has shown benefits, including social, emotional, and cultural support.⁵²⁻⁵⁵ Various models of peer support exist, with the potential to improve self-efficacy, positive mood, understanding of self-care, and perception of social support as well as improve health-related outcomes.⁵⁶ People living with chronic disease are not the only ones who may require this support; family members, caregivers, and partners of people with diabetes can experience diabetes distress.^{29,57,58} Diabetes care and education specialists play a critical role in this process by assessing individuals' support network, reinforcing the importance of this aspect of diabetes self-care, and providing training for peers and community health workers to promote accurate and appropriate messages.

Healthy Eating

Healthy eating refers to "a pattern of eating a wide variety of high quality, nutritionally-dense foods in quantities that promote optimal health and wellness."⁵⁹ The behaviors surrounding when to eat, what to eat, and how much to eat are influenced by a complex set of factors, including food and cultural preferences, food security, health beliefs, and eating habits.⁶⁰ This complexity intensifies with the additions of dynamic nutrition recommendations, health literacy challenges, varied wellness goals, and changing health status. Consequently, customization of meal plans and eating patterns based on age, activity level, health status, food preferences, and medical and nutritional management of multiple conditions among other factors becomes vital to effective behavior change.

Behaviors That Contribute to Healthier Outcomes

Develop and Use a Personalized Meal Plan

Both the Academy of Nutrition and Dietetics (AND) and the American Diabetes Association (ADA) support an evidence-based approach to individualized meal planning.^{59,61-63} In partnership with registered dietitian-nutritionists, individuals with diabetes can develop meal plans that focus on macronutrient quality, healthy eating patterns, metabolic goals, and personal food preferences.⁵⁹ Accordingly, the ADA Standards of Medical Care in

Diabetes-2020 identifies diabetes-specific medical nutrition therapy as an essential component of an overall diabetes plan.⁵⁹ Strategies such as carbohydrate counting, the plate method, and weight management exchange lists have been effective in achieving cardiometabolic and weight management goals.^{64,65} These techniques empower people to make choices based on their food preferences while maintaining an energy intake and macronutrient composition focused on their metabolic and health goals.

Establish Healthy Eating Patterns

As defined by the Dietary Guidelines 2015-2020, a healthy eating pattern contains an assortment of colorful vegetables, fruits, whole grains, low-fat dairy, a variety of protein sources, and oils while minimizing sodium, added sugars, saturated fat, and trans fat.⁶² Through discussion and problem-solving with their health care team, individuals with diabetes and related conditions can learn how to integrate healthy and safe eating patterns into their daily lives, which often requires small increments of change to achieve sustainability.⁶³

Measure Portions and Monitor Intake

Tracking the amount of food and beverages consumed for total calories, as well as for individual nutrients such as carbohydrates, plays a role in achieving weight and wellness goals. Individuals can use scales, measuring cups, apps that evaluate photo images of meals, and household measurements or hands to estimate amounts⁶⁶ and receive feedback on how their portions compare to established serving sizes. When health goals include weight loss, portion-controlled eating plans can improve weight, LDL-cholesterol, and A1C.⁶⁷ Additionally, research shows that self-monitoring of intake may predict dietary change over the long term.⁶⁸ Combining measurement and monitoring (Monitoring) within the behavior of Healthy Eating leads to Problem Solving amounts and types of food and beverages to consume to meet personalized plans.

Understand and Use the Nutrition Facts Label

Comprehension of the nutrition facts label and associated *health literacy and numeracy* skills are essential to the behavior of Healthy Eating for management of pre-diabetes, diabetes, and cardiometabolic conditions. The ability to read the nutrition facts label and calculate portions (Healthy Eating) can promote healthy eating and

drinking decisions that lower cardiovascular risk, improve glycemia (Reducing Risk), and aid in decision-making during special circumstances, such as restaurant dining or special occasions (Problem Solving).

Being Active

Being Active is inclusive of all types, durations, and intensities of daily physical movement, which equates to bouts of aerobic or resistance exercise training (structured or planned “exercise”) as well as unstructured activities. The benefits of regular physical activity on cardiometabolic health are widely known.⁶⁹

With a few exceptions, recommendations for physical participation are similar for individuals with and without diabetes. Most people with diabetes can safely begin physical activity that is no more vigorous than their usual activities of daily living without a medical checkup, which removes some barriers to their participation in increased activity.⁷⁰ When an individual with diabetes and higher cardiometabolic risk is unaccustomed to vigorous physical activity, guidelines suggest obtaining medical clearance and possibly preparticipation exercise stress testing. Comorbid health issues may require individualization of physical activity choices (eg, avoidance of weight-bearing physical activity with unhealed plantar ulcers), which serves as an opportunity for shared decision-making (Problem Solving) within the health care team. In collaboration with individuals with diabetes and related conditions, diabetes care and education specialists can provide this individualized assessment, monitor activity levels (Monitoring) as vital signs, and tackle barriers to encourage physical activity in daily lifestyles.

Behaviors That Contribute to Healthier Outcomes

The behavior changes that contribute to healthier outcomes include aerobic exercise, resistance and balance training, engaging in unstructured or daily living activities, and decreasing the amount of time spent sitting.⁷⁰⁻⁷⁵ The diabetes care and education specialist can have a particular impact in helping people identify and address barriers.

Address Barriers

Because frequent and consistent physical activity often requires sustained behavior change, combatting potential barriers with appropriate strategies and goals is of significant importance.^{76,77} Most adults discontinue

regular physical activity due to a perceived lack of time, injuries, inappropriate starting intensity, and a lack of enjoyment. Other barriers may relate to the environment, such as a lack of safe places for physical activity^{78,79}; social factors, such as a lack of social support for regular physical activity⁸⁰; and work or home situations that lead to more sedentary behaviors.

Diabetes and its related conditions present additional challenges to participation. For instance, proliferative retinopathy requires limitation of activities that cause rapid blood pressure swings, cardiac autonomic neuropathy requires extensive warm-up and cool-down phases, peripheral neuropathy requires frequent foot inspections and possibly limiting weight-bearing activity, and peripheral vascular disease requires limiting intensity to a tolerable pain threshold.⁸¹ Individuals with diabetes, especially those who use insulin or insulin secretagogues, report fear of hypoglycemia as a significant barrier.⁸² In collaboration with the health care team, people living with diabetes can develop strategies to reduce risk for or avoid hypoglycemia, such as reductions in insulin or medication prior to physical activity and inclusion of rapid-acting carbohydrate prior to and during activity (Reducing Risk and Problem Solving). Additionally, a lack of self-efficacy or self-esteem appears related to being active,⁸³ which links to the need for Healthy Coping.

Taking Medication

Medications remain an essential component in the prevention and management of chronic disease. Insufficient treatment interventions, therapeutic inertia, and/or skipping/missing medication doses⁸⁴⁻⁸⁶ continue to be barriers to reaching therapeutic goals and contribute to higher health care costs, adverse outcomes, and inferior quality of life for persons with chronic disease.^{86,87} Medication-taking behaviors include following the day-to-day prescribed treatment with respect to timing, dosage, and frequency as well as continuing treatment for the prescribed duration.⁸⁸ The reasons for not taking medications as prescribed are multifactorial.^{86,89} Given that diabetes has been recognized as a multisystem disorder with several associated comorbidities,^{90,91} treatment follows a multifaceted and individualized approach that includes cardiovascular risk mitigation.⁹⁰

Advances in the scientific understanding of diabetes have resulted in a spectrum of new oral and injectable agents targeting multiple disease mechanisms.⁹¹⁻⁹⁴ The focus has

shifted from solely an A1C reduction to a more comprehensive approach that includes consideration of time-in-range (TIR), cardiovascular disease prevention, and quality-of-life measures.⁹⁵⁻⁹⁷ Adding the dynamics of an aging population, changing demographics, social and environmental influences, access to health care, predictive medicine, and technological innovation further increases the complexities of treatment approaches and the need for both individualization and coordination of the care plan.^{85,93,98-101}

Behaviors That Contribute to Healthier Outcomes

Keep a Current, Accurate Medication List and History

A medication list provides valuable information for the individual managing health conditions and the other members of the health care team. An accurate and complete list that includes complementary therapies and over-the-counter medications can lead to collaborative discussions to address optimal selection of agents, possible conversion to newer agents with additional health benefits, deprescribing as needed, and avoidance of prescribing medications already determined to be ineffective or to cause adverse effects for this individual (Problem Solving). The medication list also serves as a mechanism to communicate medication changes within the health care team and prevent medication discrepancies from occurring, especially among various health care sites.¹⁰² Medication reconciliation among the members of the health care team, especially the individual taking the medication, may reduce the frequency of hospitalizations and emergency room visits.¹⁰³

Fill the Prescription

Filling the initial prescription, having support such as reminder prompts, and having uninterrupted and convenient prescriptions promote medication-taking behavior.⁸⁴ It can be challenging to navigate the health care system to fill a prescription¹⁰⁴; however, an understanding of the roles on the health care team and influencers of cost and coverage can facilitate the process. For instance, a person with diabetes may find that the newly prescribed medication is too expensive to fill. This problem involves Taking Medication, but it also necessitates active discussion and collaboration among members of the health care team to examine the benefits and costs for this particular individual. These factors may include (1) whether the medication's value outweighs its cost due to additional

cardiometabolic benefits (Reducing Risk), (2) whether the medication results in a noticeable improvement in this individual's blood glucose level (Monitoring), and (3) whether the benefits of adding another medication to the treatment plan offset potential financial and emotional stressors (Healthy Coping).

Take Medication as Prescribed and at the Right Time

To maximize benefits and minimize side effects, individuals may need to take medications at specific times, in relation to food, or in response to blood glucose levels. Medications may also require appropriate spacing with or from other meds. Nonoral medications (eg, injectables or inhaled insulin) or medication delivery systems (eg, insulin pumps and closed loop systems) have additional requirements to maximize effectiveness and safe use. When actual use of medications differs from prescribed plans, the members of the health care team can conduct further Problem Solving.

Share Medication Beliefs and Concerns

Health and cultural beliefs about medications are important discussion points when deciding on appropriate therapeutic options and navigating health plans for coverage for individuals with diabetes and related conditions.⁹² Taking medication as prescribed is more likely when individuals perceive medication is efficacious, namely, when they see or feel that the medication has brought a positive and immediate outcome.¹⁰⁵ Consequently, individuals may discontinue therapy when they do not experience a noticeable change. Similarly, even a single episode of hypoglycemia can affect medication-taking behavior.¹⁰⁶ As active listeners, diabetes care and education specialists can be vital to eliciting these conversations with individuals with diabetes and related conditions.¹⁰⁷ This collaboration can then lead to evaluating and addressing person-centered medication concerns, such as side effects, efficacy, cost, personal preference and lifestyle, acceptable risk of hypoglycemia, weight goals, and appropriate complexity of plan.⁹² Shared decision making (Problem Solving) around medication use enhances engagement, promotes prevention and risk reduction, and improves outcomes.¹⁰⁸⁻¹¹⁰

Monitoring

Monitoring has expanded beyond self-monitoring of blood glucose to include monitoring of blood pressure,

activity, nutritional intake, weight, medication, feet/skin, mood, sleep, symptoms like shortness of breath, and other aspects of self-care. Although monitoring can still include the use of paper and pencil to record data, new methods of data collection are available that enable individuals to more easily record data¹¹¹ and are less vulnerable to misreporting.¹¹² Furthermore, tracking lifestyle data with notes adds context to metabolic data, aiding in interpretation and more informed decision-making. With connected devices and other technologies, remote patient monitoring and virtual collaborative care is possible. Encounters are no longer time-bound; these encounters can be data-driven, and with the right timing and appropriate touchpoints, they can be influential in behavior change and in building self-management capability. Continuous glucose monitoring¹¹³ has transformed intermittent monitoring to monitoring in real time, providing insight into measures such as TIR and glucose management indicator in its continuous data delivery.

The behavior of Monitoring acts as a springboard into the other 6 self-care behaviors; the behavior itself produces data, and knowing how to use these data supports change. As an example, glucose monitoring may reveal episodes of hypoglycemia. When the data are collected (Monitoring) and shared with the health care team, they fuel discussion to find solutions (Problem Solving). Solutions may span multiple self-care behaviors, such as adjusting the time of physical activity to after a meal (Being Active), having a snack with carbohydrate (Healthy Eating), transitioning from a sulfonylurea to a GLP-1 RA (Taking Medication), maintaining a positive outlook when results are out of range (Healthy Coping), and carrying glucose gel as a precautionary measure (Reducing Risk).

Behaviors That Contribute to Healthier Outcomes

Track Appropriate and Accurate Information

The self-care behavior of Monitoring for diabetes and cardiometabolic conditions includes both metabolic and lifestyle tracking.¹¹⁴ With the ability to track more variables, the health care team must agree on which information to track and evaluate and the frequency with which to track these measures to prevent data overload and decrease time burdens associated with interpretation. Devices may require calibrations, maintenance, special instructions, and other care to capture data accurately.

Maintain and Share Organized Records

The presentation of data, whether done manually or automatically, must make sense to the individual(s) using the data. Organized records simplify interpretation and improve goal setting and shared decision-making among the members of the health care team, which can equate to medication adjustments between visits (Reducing Risks and Problem Solving).¹¹⁵ Technology can facilitate monitoring, record-keeping, and sharing data within the health care team.

Identify Trends

Structured self-monitoring of blood glucose, such as obtaining a 7-point profile of blood glucose values at fasting, preprandial and 2 hours postprandial at each meal, and bedtime, provides an opportunity to identify trends over consecutive days and improve outcomes.¹¹⁶ The revolutionary change from self-monitoring of blood glucose to continuous glucose monitoring has also underscored the value of trending data and the impact on quality of life.¹¹⁷ Closer examination of multiple data points can yield more meaningful information than an isolated value; the identification of these patterns further spurs behavior change. This overarching approach (rather than single values) may also help relieve the emotional triggers, such as pride, embarrassment, disappointment, or anger, associated with unexpected values (Healthy Coping).

Be Empowered and Engaged

The ability to see cause and effect through monitoring makes it an effective motivation tool. Engagement in self-monitoring can result in clinical improvements, such as improved blood pressure.¹¹⁸ Patient-generated health data also have the potential of improving safety, quality, care coordination, and shared decision-making.¹¹⁹ Individuals living with diabetes and related conditions actively contribute to data creation and gathering in Monitoring, empowering people to share their valuable expertise in self-care.

Reducing Risks

Reducing risk refers to identifying risks and implementing behaviors to minimize and/or prevent complications or adverse outcomes. These include hypoglycemia, hyperglycemia, diabetic ketoacidosis, hyperosmolar hyperglycemic state, retinopathy, nephropathy, neuropathy, and cardiovascular complications.¹²⁰ The ADA's Standards of Medical Care in Diabetes provides standards and evidence to promote health and decrease risk for people with diabetes.¹²¹

Behaviors That Contribute to Healthier Outcomes

Act Early

Awareness of prediabetes is positively associated with engaging in risk reduction behaviors.¹²² The National Diabetes Prevention Program provides an evidence-based approach focused on preventing or delaying the onset of type 2 diabetes for those with prediabetes or risk factors for diabetes.^{123,124} This program, focused on lifestyle interventions, has shown clinically meaningful cardiometabolic health improvements related to weight, A1C, fasting blood glucose, systolic and diastolic blood pressure, and total and HDL-cholesterol.^{125,126} These lifestyle modifications tie to the self-care behaviors of Healthy Eating and Being Active, in addition to Taking Medications.

Participate in DSMES

Research has demonstrated that participating in DSMES can lower A1C by as much as 1%¹²⁷ and that A1C improvements are associated with a decrease in microvascular and potentially macrovascular complications.¹²⁸⁻¹³⁰ Research also reveals that this participation has decreased acute care visits, hospitalizations, and readmissions,¹³¹ which contributes to cost savings within the health care system. Participation in DSMES can increase utilization of primary care and preventive services, such as laboratory testing, eye and dental exams, and screenings for complications, aligning with best practice treatment measures over time.¹³²⁻¹³⁴

Aim for Adequate Sleep

People with diabetes are more likely to sleep poorly due to sleep apnea, restless legs syndrome, peripheral neuropathy, depression, hypoglycemia, and hyperglycemia. This lack of or poor quality of sleep contributes to elevated blood glucose and A1C levels, weight gain, and an increased risk of heart disease and obesity. Lifestyle measures and other treatments are available to promote sleep and lower the risk of complications resulting from inadequate rest.¹³⁵

Plan and Do

Individualized risk reduction practices, such as receiving vaccines for flu, pneumonia, and hepatitis B and participating in behavior change programs related to tobacco use, have the potential to improve health across populations.¹³⁶ A seemingly simple recommendation to obtain a dilated eye exam requires individuals to complete several

Table 1

Diabetes Self-Management Education and Support Core Outcome Measures: Healthy Coping

Outcomes Measurement Process				
Measurement/Assessment		Monitoring		Management
DSMES Core Outcome Measures (Diabetes Self-Care Behaviors)	Immediate Outcome Learning and Barrier Resolution	Intermediate Outcome/Behavior	Recommended Interval Between Measurement	Outcomes Information Used to Drive Decision-Making and the Delivery of Care
Healthy Coping - Internal and external motivators - Benefits of solution-focused problem-solving - Active self-management - Value of nurturing support system (peers, online, family, friends) - Individual empowerment - Role as partner with other members of health care team	Knowledge - Goal setting - Problem-solving - Coping strategies - Self-efficacy Barriers - Physical - Financial - Emotional - Competing priorities - Lack of support network - Psychosocial distress including diabetes distress - Cognitive including mental health disorders	Measures - Depression score - Stress level - Quality of life (perceived self-efficacy, perceived disease severity, perceived interference of chronic disease) - Functional measurement - Treatment self-efficacy - Level of empowerment - Absenteeism - Presence of support Methods of Measurement - Self-report - Skills, Confidence, and Preparedness Index (SCPI) - Problem Areas in Diabetes (P.A.I.D) - Quality of Life (QOL) tools, such as Short Form-36 (SF-36) or Short Form-12 (SF-12) with Appraisal of Diabetes Scale (ADS) - Depression/diabetes distress tools, such as Diabetes Distress Scales (DDS), Parents-DDS, Partners-DDS, T1-DDS or DDS; - Beck-Depression-Inventory (BDI); Patient Health Questionnaire-9 (PHQ-9) - Cognitive impairment tools, such as Saint Louis University Mental Status (SLUMS); Mini-Mental Status Exam (MMSE)	Learning Outcomes - Evaluated with each encounter Behavioral Outcomes - Baseline 2 to 4 wk - Every 3 to 6 mo - During transition periods, such as development of complications or life-cycle changes	Behavior (infrequent or inconsistent self-monitoring blood glucose and medication taking): The diabetes care and education specialist notes a change in Mark's blood glucose records. Mark appears frustrated with himself for not performing self-monitoring of blood glucose and not taking medication as recommended. Barrier identification (depression, feeling overwhelmed): Mark explains feeling overwhelmed and depressed most of the time. The diabetes care and education specialist acknowledges his feelings and asks if Mark would be willing to take a short survey to evaluate these symptoms. Mark agrees to take the PHQ-9, which yields a result of mild to moderate depression. Barrier resolution (follow-up for depression recommended): The diabetes care and education specialist and Mark discuss the PHQ-9 results. Mark agrees to see a behavioral health specialist and primary care provider for treatment. Behavior change (increased/more consistent self-monitoring of blood glucose and medication taking): At the 3-mo follow-up, Mark reports a greater interest in taking charge of his health after starting an antidepressant and behavior therapy. Mark shares his monitoring records and is proud of the changes he has made.

Abbreviation: DSMES, Diabetes Self-Management Education and Support.

Table 2

Diabetes Self-Management Education and Support Core Outcome Measures: Healthy Eating

Outcomes Measurement Process				
DSMES Core Outcome Measures (Diabetes Self-Care Behaviors)	Measurement/Assessment		Monitoring	Management
	Immediate Outcome Learning and Barrier Resolution	Intermediate Outcome/Behavior Change	Recommended Interval Between Measurement	Outcomes Information Used to Drive Decision-Making and the Delivery of Care
Healthy Eating	Knowledge - Effect of foods/beverages on metabolic parameters (including blood glucose, lipids, blood pressure, weight, etc.) - Sources and distribution of nutrients (nutrient-dense carbohydrates, lean proteins, healthy fats) - Eating patterns (frequency of meals, timing, portions, etc.) - Resources to assist in food choices - Macronutrient composition (quality, quantity, combination, substitutions) Skills - Meal planning - Portion awareness and management - Planning strategies (carb counting, exchanges, plate method, mindful eating) - Nutrition facts label comprehension - Special situations and problem-solving (planning, shopping, meal delivery/kits, eating away from home at work/school/restaurants) Barriers - Environmental factors - Cultural and family influences - Food and health beliefs - Financial (food security) - Cognitive - Health literacy and numeracy - Emotional - Meal pattern sustainability	Measures - Types of food choices - Amounts consumed - Timing of meals and snacks - Alcohol (with or without food, amount, frequency) - Fluids (adequate hydration) - Effect of food/beverages on metabolic parameters - Progress toward goal achievement Methods of Measurement - Observation - Self-report (24-h recall, typical day, food frequency, food diaries) - Monitoring tools with associated records - Goal setting	Learning Outcomes - Evaluate with each encounter - Ongoing self-evaluation and adjustments with life-cycle events and secondary diseases Behavioral Outcomes - Baseline 2 to 4 wk - Every 3 to 6 mo - Annual follow-ups - When lifestyle or health status changes	Behavior (inconsistent food intake): Suzy shares her food and blood glucose records. The diabetes care and education specialist acknowledges Suzy's self-monitoring efforts and reviews her records. They discuss eating behaviors, such as skipped meals and overeating. Barrier identification (ineffective problem-solving): Suzy does not plan for eating meals/snacks when away from home (for work or school); risks include food availability and timing issues. Barrier resolution (increased knowledge regarding importance of meal planning): The diabetes care and education specialist and Suzy work together to identify potential healthy eating patterns, identify Susie's strengths and set a realistic goal to apply those strengths for change. Suzy becomes more aware of her eating when away from home. She considers options to improve meal planning for meals/snacks at home and away. She reports increased energy and positive feeling of successful self-management when she participates in these behaviors. Behavior change (consistent food intake, improved blood glucose, cardiometabolic parameters, weight management): Suzy now plans, shops, and packs meals/snacks in advance. She also reviews menus of other food options for purchase. Blood glucose levels and labs more consistently in the target range are evident at 6-mo follow up.

Table 3

Diabetes Self-Management Education and Support Core Outcome Measures: Being Active

Outcomes Measurement Process				
DSMES Core Outcome Measures (Diabetes Self-Care Behaviors)	Measurement /Assessment		Monitoring	Management
	Immediate Outcome Learning and Barrier Resolution	Intermediate Outcome/Behavior	Recommended Interval Between Measurement	Outcomes Information Used to Drive Decision-Making and the Delivery of Care
Being Active	Knowledge - Planned exercise (type, duration, intensity, frequency, progression) - Daily movement - Breaking up sedentary time - Safety precautions, such as obtaining preparticipation medical clearance and/or exercise stress testing prior to unaccustomed vigorous activity - Special considerations, such as appropriate footwear Skills - Appropriate daily movement and physical activity plan - Adjustment of activity with food and medication to maintain glycemic balance - Monitoring of cardiometabolic parameters, data stream, and feedback Barriers - Physical (health conditions, injuries) - Perceived lack of time - Environment, facilities - Fear (hypoglycemia) - Self-efficacy - Lack of enjoyment - Lack of social support	Measures - Type, frequency, duration, and intensity of planned activities - Daily movement - Progress toward goal achievement - Quality of life, health improvement Methods of Measurement - Self-report - Goal setting - Monitoring tools and their associated records, including digital health tracking and wearable technologies - Quality of life and health assessments - Exercise vital sign (EVS) to evaluate whether weekly goals for physical activity have been met - On average, how many days per week do you engage in moderate to strenuous exercise (like a brisk walk)? - On average, how many minutes do you engage in exercise at this level? - Physical Activity Vital Sign (PAVS) when individual is physically active for at least 30 min per day - How many days during the past week have you performed physical activity where your heart beats faster and your breathing is harder than normal for 30 min or more? - How many days in a typical week do you perform activity such as this?	Learning Outcomes - Evaluate with each encounter Behavioral Outcomes - Baseline 2 to 4 wk - Every 3 to 6 mo	Behavior (lack of physical activity): When prompted by her diabetes care and education specialist, Arya discusses/shares that she's had little success in her goal of increasing physical activity over the last 6 wk. Barrier identification (environment): The diabetes care and education specialist and Arya discuss barriers and discover previous success at an exercise facility years ago. Arya currently does not feel comfortable attending due to her increased weight and perceived body image. Behavior resolution (environment changed): Through discussion and problem-solving with the diabetes care and education specialist, Arya agreed to try a women's-only exercise club near work that was convenient and nonthreatening. Behavior change (increased activity): At 1-mo follow-up, Arya reports performing aerobic exercise (planned) 3 to 5 days per week. She increased her total daily movement and already lost 5 pounds. Behavior (lack of physical activity): At her virtual appointment with her diabetes care and education specialist, Arya shares her desire to be more active but is unsure how to get started. Barrier Identification (financial): The diabetes care and education specialist and Arya discuss options. Arya discloses that she cannot afford a gym membership but is willing to purchase a wearable device. The diabetes care and education specialist and Arya review basics of a wearable device and strategies to stay engaged. Barrier resolution (try wearable): Arya purchases a wearable device for activity tracking and joins a weekly virtual contest on the associated mobile app. She wants to win the weekly contest by getting the most steps and agrees on an initial goal of 4000 steps per day on most days of the week. Behavior change (increase activity): At a 1-mo follow-up, Arya shares her activity tracker report with the diabetes care and education specialist, which reveals an average step count of 3000 to 5000 on most days of the week. The diabetes care and education specialist acknowledges Arya for her commitment to physical activity and encourages Arya to increase her goal. Arya feels confident in setting a goal for 7000 steps for 5 days per week within the next month.

Table 4

Diabetes Self-Management Education and Support Core Outcome Measures: Taking Medication

Outcomes Measurement Process				
DSMES Core Outcome Measures (Diabetes Self-Care Behaviors)	Measurement/Assessment		Monitoring	Management
	Immediate Outcome Learning and Barrier Resolution	Intermediate Outcome/Behavior	Recommended Interval Between Measurement	Outcomes Information Used to Drive Decision-Making and the Delivery of Care
Taking Medication	Knowledge - Name, dose, frequency, and optimal timing of medications - Medication mechanism of action - Common side effects, toxicity - Action for adverse effects - Action for missed dose - Storage, travel, safety, and disposal - Recognition of efficacy, optimal outcomes, and therapeutic goals Skill - Maintenance of a medication list - Preparation, technique, administration - Safe handling, disposal of equipment - Dose adjustment - Recognition, treatment, prevention of common adverse effects Barriers - Plan complexity (> 1 medication or dose daily) - Physical (vision or dexterity) - Financial (medication cost, copay) - Health beliefs (skeptical of benefit, worried about side effects) - Health literacy and numeracy - Cognitive (dose recall, refill initiation) - Psychological (depression, fear, or embarrassment) - Change in schedule or work status	Measures - Medication taking - Prescription filling - Dose accuracy - Glycemic trends - Metabolic trends - Emergency department and hospital utilization - Weight change Methods of Measurement - Self-report and medication records - Review of pharmacy refill history - Pill count - Return demonstration, (observation, role-playing) - Labs (A1C, total cholesterol, LDL-cholesterol, etc) - Monitoring tools with associated records (eg, records for blood glucose, blood pressure, weight, medication use, etc)	Learning Outcomes - Evaluated with each encounter Behavioral Outcomes - Baseline 2 to 4 wk (may be earlier in cases such as new to insulin therapy) - Every 3 to 6 mo or if medication concerns are suspected	Behavior (adapting to new medication/dose): The diabetes scare and education specialist and Kim review his current medication list. The diabetes specialist identifies a new dose of medication listed in the electronic health record that does not match with Kim's list. The diabetes care and education specialist asks about the new dose, and Kim reports that he has not picked it up at the pharmacy. Barrier identification (lack of understanding): The diabetes care and education specialist and Kim review the medication list and discuss the new dose. After asking a few questions, the diabetes care and education specialist uncovered that Kim did not understand the dose change instructions at the last visit. The diabetes care and education specialist and Kim update his medication list together and discuss his concerns about the new medication regimen. Barrier resolution (increased knowledge of medication plan): By the end of the visit, Kim can accurately describe the new plan and reports feeling more comfortable with the change. They outline a plan for Kim to obtain the new dose of medication. Behavior change (medication-taking behavior): The diabetes care and education specialist contacts Kim through an electronic message in a web portal that links to the electronic health record. Kim reports taking the new dose of medication as prescribed. Kim uploads his blood glucose records to the portal for review by the diabetes care and education specialist and other members of the health care team. The diabetes care and education specialist replied with an electronic message to acknowledge Kim's efforts toward the increase in time-in-range and encouraged him to continue with his medication-taking behavior.

Table 5

Diabetes Self-Management Education and Support Core Outcome Measures: Monitoring

Outcomes Measurement Process				
DSMES Core Outcome Measures (Diabetes Self-Care Behaviors)	Measurement/Assessment		Monitoring	Management
	Immediate Outcome Learning and Barrier Resolution	Intermediate Outcome/Behavior	Recommended Interval Between Measurement	Outcomes Information Used to Drive Decision-Making and the Delivery of Care
Monitoring	Knowledge - Monitoring plan/schedule (structured, episodic, continuous, etc) - Appropriate lifestyle data to track - Target values - Safety issues including disposal of lancets - Use of data for decision-making - Awareness of body's symptoms (eg, blurred vision, shortness of breath) and/or physical changes (eg, teeth, skin, gums) Skills - Equipment use and technical care (blood glucose meter, continuous glucose monitor, blood pressure cuff, wearable, mobile app, etc) - Record keeping with note taking - Tracking and reporting body symptoms and physical changes - Interpretation of patient-generated health data Barriers - Physical - Financial - Cognitive - Emotional - Time - Inconvenience - Treatment burden - Health literacy and numeracy - Limited understanding of value of data and how to use it - Lack of interest/ability to use equipment and other tools for self-monitoring	Measures - Frequency of self-monitoring - Schedule of monitoring "Unscheduled" monitoring (triggered by symptoms, etc) - Number of devices/apps used to support monitoring - Blood glucose values - Time-in-range (TIR) - Glucose management indicator (GMI) - Blood pressure values - Hours of sleep - Mood status - Amount of time performing physical activity, number of steps - Medication use/insulin doses - Amount of carbohydrate consumed, meal size - Presence of notes that add context to tracked data - Presence of organized data that allow for decision-making Methods of Measurement - Monitoring tools and their associated records (log book, device memory review, printouts) - Self-report responses to questions/surveys	Learning Outcomes - Evaluate with each encounter (Review of automated data provides insights into knowledge and use of monitoring device or devices) Behavioral Outcomes - Baseline 2 to 4 wk - Every 3 to 6 mo (May do more often based on virtual care/remote monitoring program guidelines)	Behavior (minimal self-monitoring of blood glucose [SMBG]): Malik shares his blood glucose records with his diabetes care and education specialist. The diabetes care and education specialist identifies sporadic monitoring on weekdays. Barrier identification (treatment burden): The diabetes care and education specialist acknowledges Malik's record keeping and asks about the gaps. Malik reports that he avoids checking blood glucose at work. Barrier resolution (use of continuous glucose monitoring [CGM]): The diabetes care and education specialist reviews other options to capture blood glucose with Malik. Malik is curious about the continuous glucose monitor and agrees to try it. Behavior change (obtaining blood glucose data through SMBG and CGM): After instruction on the equipment and calibrations, Malik felt ready to wear a CGM for a trial of 1 wk. He continues to prefer this method of monitoring blood glucose and provides these records to his health care team for shared decision-making regarding his insulin plan. Behavior (monitoring blood pressure): While at his appointment with the diabetes care and education specialist, Malik acknowledges the importance of managing his blood pressure and feels frustrated that he can only see blood pressure measurements during office visits. Barrier identification (financial/health literacy): The diabetes care and education specialist and Malik discuss the possibility of taking blood pressure measurements at home. Malik is concerned about the cost of a blood pressure monitor. Barrier resolution (health insurance navigation and equipment training): The diabetes care and education specialist works with Malik to help him obtain a blood pressure monitor through his health insurance plan. The diabetes care and education specialist teaches Malik how to use it to produce accurate results. Behavior change (additional monitoring for hypertension): Malik now tracks his blood pressure outside of office visits and generates a report to share with his health care team for collaboration and treatment adjustments.

Table 6

Diabetes Self-Management Education and Support Core Outcome Measures: Reducing Risks

Outcomes Measurement Process				
DSMES Core Outcome Measures (Diabetes Self-Care Behaviors)	Measurement/Assessment		Monitoring	Management
	Immediate Outcome Learning and Barrier Resolution	Intermediate Outcome/Behavior	Recommended Interval Between Measurement	Outcomes Information Used to Drive Decision-Making and the Delivery of Care
Reducing Risks	Knowledge • Safety (sick day plan, driving/machine operation precautions, emergency preparedness) • Standards of care • Therapeutic goals • Symptoms that require attention or follow-up (hypoglycemia, hyperglycemia, rapid weight fluctuation, stroke, heart attack, bleeding gums, vision changes, skin changes) • How to decrease risks/prevent harm (pregnancy counseling, smoking cessation, etc) Skills • Planning • Monitoring of blood glucose (self, continuous) • Maintaining personal care record • Performing self-foot exam • Performing self-skin exam • Self-monitoring of blood pressure • Use of health apps and web portals • Ability to adjust food, medication, and activity (to increase the amount of time glucose is in range) • Recognition of concerning symptoms or changes in health • Ability to determine when health requires care from health care team (emergency vs nonemergency)	Measures • Glycemic trends • Frequency of low or high blood glucose • Frequency of contact with health care provider for problem resolution • Missed days from work, school, or related activities • Number of visits to the emergency department or hospitalizations • A1C • Lipids • Blood pressure • Kidney tests (urine albumin excretion and serum creatinine for eGFR, urine albumin creatinine ratio) • Weight and body mass index (BMI) • Scheduled vs attended visits with health care team • Dilated eye exam • Immunization status (flu vaccine, pneumonia vaccine, hepatitis B) • Screening for hearing loss • Dental exam • Sleep study • Smoking status • Frequency of foot self-exam • Comprehensive foot exam • Screening for sexual dysfunction • Neuropathy • Aspirin therapy	Learning Outcomes • Evaluated with each encounter Behavioral Outcomes • Baseline • 2 to 4 wk • Every 3 to 6 mo • Annual follow-ups	Behavior (frequent hypoglycemia): During her visit with the diabetes care and education specialist, Rida reports 5 episodes of hypoglycemia in the last month. Rida notes that these episodes occurred after doing extra work in her vegetable garden. Barrier identification (knowledge deficit): After discussion with the diabetes care and education specialist about hypoglycemia, Rida realized she had not considered gardening as physical activity. Additionally, she had not realized that her medication (sulfonylurea) increased the likelihood of hypoglycemia. Barrier resolution (information provided): The diabetes care and education specialist reviewed signs, symptoms, and treatment of hypoglycemia. The diabetes care and education specialist and Rida discussed scenarios with increased risk of hypoglycemia and then problem-solved how to balance food, activity, and medication to reduce risk for hypoglycemia. Behavior change (decrease in hypoglycemia): During her 1-mo follow up, Rida reported only 1 episode of hypoglycemia. She was able to detect signs ("starting to feel shaky") after taking a walk with a neighbor. Rida reports now carrying a source of glucose with her, treating low values with approximately 15 g of carbohydrate, and rechecking blood glucose until a return to a safe blood glucose level. Behavior (no eye exam): The diabetes care and education specialist asked Rida about her preventive exams this year. Rida reports that her most recent dilated eye exam was 3 y ago. Barrier identification (knowledge deficit): Rida does not believe she needs a dilated eye exam because she is not experiencing visual disturbances. She is content with the magnifying glasses she uses.

(continued)

Table 6

(continued)

	Outcomes Measurement Process			
	Measurement/Assessment	Monitoring	Management	
DSMES Core Outcome Measures (Diabetes Self-Care Behaviors)	Immediate Outcome Learning and Barrier Resolution	Intermediate Outcome/Behavior	Recommended Interval Between Measurement	Outcomes Information Used to Drive Decision-Making and the Delivery of Care
	Barriers - Financial (lack of personal resources; insurance barriers such as high deductible, underinsured, step therapy requirements; insufficient monitoring supplies; food insecurity) - Unawareness of disease process and its seriousness - Lack of access to diabetes self-management education services or health care providers - Therapeutic inertia - Physical (hypoglycemia unawareness) - Cognitive - Emotional - Lack of self-efficacy and coping strategies - Poor support network, including lack of rapport with provider - Perceived lack of time	- Frequency of medication adjustment - Sick day plan Methods of Measurement - Self-report - Chart or exam code audit - Review of monitoring records - Demonstration of self-care activities		Barrier resolution (knowledge of importance provided): Rida reports that she had not known that her eyes were affected by diabetes. The diabetes care and education specialist and Rida discussed the rationale of preventive dilated eye exams and recommended frequency. Rida agreed to make an appointment after confirming that this was a covered benefit under her insurance plan. Behavior change (dilated eye exam done): Rida sent a message to her diabetes care and education specialist through the portal to let the diabetes care and education specialist know that she had completed her dilated eye exam. Rida obtained a copy of her report and shared it with the rest of her health care team through the portal.

Table 7

Diabetes Self-Management Education and Support Core Outcome Measures: Problem Solving

Outcome Measurement Process			
Measurement/Assessment		Monitoring	Management
DSMES Core Outcome Measures (Diabetes Self-Care Behaviors)	Immediate outcome Learning Barrier Resolution	Intermediate Outcome/Behavior	Outcomes Information Used to Drive Decision-Making and the Delivery of Care
Problem Solving • Knowledge • Complexity and challenges of diabetes • Changes in diabetes throughout the life cycle • Changes in diabetes as it progresses • Relevant diabetes self-management education and support knowledge items (see other behaviors) • Skills • Relevant diabetes self-management education and support skill items • Ability to recognize/identify problem • Ability to generate potential solutions • Ability to transfer past experience(s) • Ability to finalize solution • Ability to measure/monitor results	Measures • Glycemic trends including time-in-range (TIR) and glucose management indicator (GMI) • A1C • Other health indicators (weight, blood pressure, etc) • Frequency of phone calls/visits to provider • Quality of life indicators (missed days of work/school, frequency of hypoglycemia, etc) • Confidence level (in situational problem-solving) • Progress toward goal achievement • Level of diabetes distress • Frequency of acute complications • Problem Areas in Diabetes (PAID) score • Frequency of medication adjustment Methods of Measurement • Self-report • Return demonstration/teach-back • Goal setting • Monitoring tools and associated records (data from meter, continuous glucose monitor, device, app, lab) • Health Problem Solving Scale (HPSS) • Summary of Diabetes Self-Care Activities scale (SDSCA)	Learning outcomes • Evaluated with each encounter Behavioral outcomes • Baseline • 2 to 4 wk • Every 3 to 6 mo • When perceived problems arise	Behavior (lack of blood glucose monitoring): Haris reports frequent hypoglycemia in the past month but did not have any blood glucose records to share. The diabetes care and education specialist probed further to uncover that all of these episodes of hypoglycemia occurred at work. Barrier identification (coping): The diabetes care and education specialist and Haris review his workday, medication-taking behaviors, eating, and activity habits. Haris pinpoints that he has had an increased workload and sometimes skips lunch. Barrier resolution (problem solving): The diabetes care and education specialist and Haris work together to clarify the problem and identify potential solutions. Haris does not feel comfortable approaching his supervisor about taking lunch. However, he is willing to talk with human resources to identify appropriate breaks at work. The diabetes care and education specialist and Haris discuss the plan and role-play the discussion. Behavior change (decrease in hypoglycemia): After confirming his allowed breaks, Haris chooses to make time for lunch at work. He reports 1 episode of hypoglycemia in the past month, unrelated to work. Haris agrees to share his blood glucose records with his diabetes care and education specialist for additional follow-up.

steps and make decisions. In this example, obtaining a dilated eye exam requires Problem Solving to determine which provider to use and at what cost, schedule the appointment, handle the logistics of getting to the appointment, and then actively participate in the visit (Reducing Risk). Being able to plan an activity and divide it into achievable tasks fosters success in this behavior.

Engage in Health

Information gathered from self-monitoring (Monitoring) or from labs ordered by a provider (eg, A1C, lipids, or kidney function tests) can serve as the basis for interest in improving self-care behaviors. Using data from self-monitoring of blood pressure and of blood glucose can contribute to effective self-management, lowering diabetes and cardiometabolic complications.^{90,137,138} Individuals become engaged in their health when they take a more active role, such as preparing questions in advance for their appointments, sharing their health records, or inquiring about their lab results. Active participation in health also relies on an individual's skills in Healthy Coping, to be able to (1) acknowledge the value in preventing health problems that are not yet tangible and (2) recognize the power of individual behavior to change health outcomes.

Problem Solving

Problem solving is defined as “a learned behavior that includes generating a set of potential strategies for problem resolution, selecting the most appropriate strategy, applying the strategy, and evaluating the effectiveness of the strategy.”⁵ It is an essential skill for effectively self-managing diabetes and successfully implementing desired behaviors.^{5,139} In fact, DSMES uses problem-solving as a strategy to facilitate goal setting, goal achievement, and skill attainment. The National Standards for Diabetes Self-Management Education and Support¹⁶ recommend curricula designs that address decision-making and problem-solving skills. At the most basic level, problem-solving is a process that involves 3 steps: (1) Identify the problem, (2) develop alternative solutions, and (3) select, implement, and evaluate the solutions.¹⁴⁰

The use of problem-solving skills has been associated with positive clinical outcomes, specifically an improvement in A1C.^{141,142} Problem-solving models have proven effective among low-income urban^{143,144} and rural underserved populations.¹⁴⁵ The ability to problem solve is affected by self-efficacy (Healthy Coping). When individuals succeed in solving their self-identified problems,

they gain confidence in their ability to handle future challenging situations,¹⁴⁶ enhancing their self-efficacy.

Behaviors That Contribute to Healthier Outcomes

Unlike the other 6 self-care behaviors, a foundation of knowledge and skills in the other 6 behaviors is helpful for effective Problem Solving in diabetes self-management.

Ask for Clarification and Disclose Challenges

The language and terms used in health care can be confusing and difficult to understand. The Centers for Disease Control and Prevention estimate the challenge of health literacy and numeracy affects 9 out of 10 adults.¹⁴⁷ Moreover, cognitive ability is strongly associated with literacy skills; these abilities include storage of knowledge (crystallized abilities) and the ability to learn and apply new information (fluid abilities).⁵¹ Due to the complexity of problem-solving, this self-care behavior requires a sustained partnership between individuals, diabetes care and education specialists, and health care providers. This implies that individuals need to openly share their concerns and discuss their limitations, and the other members of the health care team need to listen closely and thoroughly assess individuals' knowledge, skills, and barriers; this teamwork encourages proper problem identification and effective self-management.

Participate in Shared Decision-Making and Collaborative Goal Setting

Various approaches exist to support shared decision-making, including transferring information, prioritizing decisions, and discussing the advantages and disadvantages of available choices.¹⁴⁸ Goal setting is also an essential component of collaborative care; it creates a sense of purpose while increasing positive solution-based thinking.¹⁴⁹ Research shows that goal setting is positively associated with A1C levels.^{150,151} Because self-identified problems or goals are most relevant to the individual with diabetes or related conditions, these problems are appropriate starting points for collaboration and shared decision-making among the members of the health care team.

Create an Environment That Promotes Health

Many factors, such as economic stability, employment, education, social and community context, health care systems, and neighborhoods, influence health and therefore

affect the ability of an individual to self-manage a health condition. These social determinants of health are defined as “conditions in the environments in which people live, learn, work, play, worship, and age that affect a wide range of health, functioning, and quality-of-life outcomes and risks.”¹⁵² A positive physical and social environment can reduce or eliminate barriers within each self-care behavior, facilitating the development of problem-solving skills.

Be a Lifelong Learner and Learn From Choices

An individual's health, as well as advancements in science and health care, change over time, creating the need for lifelong learning. People at risk for and with diabetes and related conditions, as well as their health care teams, benefit from being active learners who seek out information and are assertive in their pursuit to understand new information. Learning from previous choices and then revising plans based on information gained facilitates behavior change and mastery of problem-solving skills, creating a cycle of continual improvement in self-management.¹⁵³

ADCES7: Self-Care + Care Team = Optimal Outcomes

The ADCES7 model provides a plan for individuals living with diabetes and related conditions to support self-care. It also guides clinical, behavioral, and educational assessment for the health care team. The ADCES7 framework serves as a benchmark in continuous quality improvement activities in an effort to measure, monitor, and manage behavior change. Moreover, the ADCES7 framework enables individualized, comprehensive care¹⁵⁴ by health care providers using a person-centered¹⁴ and team-based approach.¹⁵⁵ Team-based care has the potential to improve satisfaction, decrease costs, lower readmission rates, and improve health.¹⁵⁶

For individuals living with diabetes and related conditions, diabetes care and education specialists are key contributors to team-based care. Using the ADCES7 framework, diabetes care and education specialists address educational, clinical, psychosocial, and behavioral needs and customize strategies²⁷ to attain “optimal health and quality of life for persons with, affected by, or at risk for diabetes and other chronic conditions.”² Their expertise spans therapy optimization, care coordination, care plan

development, and integration of new technologies to help achieve the Institute for Healthcare Improvement's Quadruple Aim to: (1) improve the health of the population, (2) enhance experiences and outcomes of individuals receiving care, (3) decrease per capita costs, and (4) improve the work life of health care providers.¹⁵⁷

Conclusion

It is the position of the Association of Diabetes Care & Education Specialists that at the cornerstone of diabetes self-management education and support, the ADCES7 is the framework for achieving behavior change that leads to effective self-management through improved behavior and clinical outcome measures. The ADCES7 Self-Care Behaviors provides a practical model that informs decision-making among individuals living with diabetes and related conditions and the members of their health care team in their shared drive for improved health and quality of life. The use of technology has transformed the approach to diabetes self-care and implementation of the ADCES7 framework. With the practical framework integrating technology, diabetes care and education specialists have the professional expertise to lead and optimize health care delivery. The ADCES7 Self-Care Behaviors demonstrate that learning, behavioral, clinical, and use of technology effectively, improves the clinical and quality of life outcomes for diabetes, cardiometabolic, related conditions, and beyond.

References

1. Association of Diabetes Care & Education Specialists. ADCES7™ Self-Care Behaviors position statement, 2014. Accessed November 2, 2020. https://www.diabeteseducator.org/docs/default-source/legacy-docs/_resources/pdf/publications/aade7_position_statement_final.pdf?sfvrsn=4
2. Association of Diabetes Care & Education Specialists. Vision for the specialty. Published 2019. Accessed November 2, 2020. www.diabeteseducator.org/Vision
3. Burke SD, Sherr D, Lipman RD. Partnering with diabetes educators to improve patient outcomes. *Diabetes Metab Syndr Obes*. 2014;7:45-53. doi:10.2147/DMSO.S40036
4. Greenwood DA, Gee PM, Fatkin KJ, Peebles M. A systematic review of reviews evaluating technology-enabled diabetes self-management education and support. *J Diabetes Sci Technol*. 2017;11(5):1015-1027. doi: 10.1177/1932296817713506
5. Mulcahy K, Maryniuk M, Peebles M, et al. Diabetes self-management education core outcomes measures. *Diabetes Educ*. 2003;29(5):768-803. doi.org/10.1177/014572170302900509
6. Peebles M, Tomky D, Mulcahy K, et al. Evolution of the American Association Diabetes Educators diabetes education

- outcomes project. *Diabetes Educ.* 2007;33(5):794-817. doi:org/10.1177/0145721707307615
7. Cornell S, Halstenson C, Miller DK. eds. *The Art and Science of Diabetes Care and Education*. 5th ed. Association of Diabetes Care & Education Specialists; 2021.
 8. Tomky D, Weaver T, Mulcahy K, Peeples M. Diabetes education outcomes: what educators are doing. *Diabetes Educ.* 2000;26(6):951-954. doi:10.1177/014572170002600607
 9. Tomky D, Cypress M, Dang D, Maryniuk M, Peyrot M, Mensing C. American Association of Diabetes Educators position statement: AADE7 Self-Care Behaviors. *Diabetes Educ.* 2008;34:445-449. doi:10.11770145721708316625
 10. Funnell M, Haas LB. National standards for diabetes self-management education programs. *Diabetes Care.* 1995;18(1):100-116. doi:10.1177/014572170302900510
 11. Mulcahy K, Maryniuk M, Peeples M, et al. Position statement: standards for outcomes measurement of DSME. *Diabetes Educ.* 2003;29(5):804-818. doi:10.2337/diacare.18.1.100
 12. Peyrot M, Peeples M, Tomky D, Charron-Prochownik D, Weaver T. Development of the American Association of Diabetes Educators diabetes self-management assessment report tool. *Diabetes Educ.* 2007;33(5):818-826. doi:10.11770145721707307614
 13. Institute of Medicine. *Crossing the Quality Chasm: A New Health System for the 21st Century*. The National Academies Press; 2001. doi:10.17226/10027
 14. Mensing C, Boucher J, Cypress M, et al. National standards for diabetes self-management education. *Diabetes Care.* 2000;23(5):682-689. doi:10.2337/diacare.23.5.682
 15. Haas L, Maryniuk M, Beck J, et al. National standards for diabetes self-management education and support. *Diabetes Care.* 2012;35(11):2393-2401. doi:10.2337/dc12-1707
 16. Beck J, Greenwood DA, Blanton L, et al. 2017 National standards for diabetes self-management education and support. *Diabetes Educ.* 2017;43(5):449-464. doi:10.2337/dci17-0025
 17. Peters A, Ahmann AJ, Battelino T, et al. Diabetes technology-continuous subcutaneous insulin infusion and continuous glucose monitoring in adults: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab.* 2016;101(11):3922-3937. doi:10.1210/jc.2016-2534
 18. Afshin A, Babalola D, McLean M, et al. Information technology and lifestyle: a systematic evaluation of internet and mobile interventions for improving diet, physical activity, obesity, tobacco, and alcohol use. *J AM Heart Assoc* 2016;5(9):e003058. doi:10.1161/JAHA.115.003058
 19. Hou C, Carter B, Hewitt J, Francisa T, Mayor S. Do mobile phone applications improve glycemic control (HbA1c) in the self-management of diabetes? A systematic review, meta-analysis, and GRADE of 14 randomized trials. *Diabetes Care.* 2016;39(11):2089-2095. doi:10.2337/dc16-0346
 20. Chiauzzi E, Rodarte C, DasMahapatra P. Patient-centered activity monitoring in the self-management of chronic health conditions. *BMC Med.* 2015;13(1):77. doi:10.1186/s12916-015-0319-2
 21. Wang J, Coleman DC, Kanter J, et al. Connecting smartphone and wearable fitness tracker data with a nationally used electronic health record system for diabetes education to facilitate behavioral goal monitoring in diabetes care: protocol for a pragmatic multi-site randomized trial. *JMIR Res Protoc.* 2018;7(4):e10009. doi:10.2196/10009
 22. Ohno-Machado L. Informatics systems for health care providers, patients, and families. *JAMA.* 2018;25(2):110. doi:10.1093/jamia/ocy001
 23. Funk M, Taylor EL. Pedometer-based walking interventions for free-living adults with type 2 diabetes: a systematic review. *Curr Diabetes Rev.* 2013;9(6):462-471. doi:10.2174/15733998113096660084
 24. Wang Y, Xue H, Huang Y, Huang L, Zhang D. A systematic review of application and effectiveness of mHealth interventions for obesity and diabetes treatment and self-management. *Adv Nutr.* 2017;8(3):449-462. doi:10.3945/an.116.014100
 25. Badawy SM, Barrera L, Sinno MG, Kaviany S, O'Dwyer LC, Kuhns LM. Text messaging and mobile phone apps as interventions to improve adherence in adolescents with chronic health conditions: a systematic review. *JMIR mHealth uHealth.* 2017;5(5):e66. doi:10.2196/mhealth.7798
 26. van Beers CA, DeVries JH, Kleijer SJ, et al. Continuous glucose monitoring for patients with type 1 diabetes and impaired awareness of hypoglycaemia (IN CONTROL): a randomised, open-label, crossover trial. *Lancet Diabetes Endocrinol.* 2016;4(11):893-902. doi:10.1016/S2213-8587(16)30193-0
 27. Powers MA, Bardsley J, Cypress M, et al. Diabetes self-management education and support in type 2 diabetes: a joint position statement of the American Diabetes Association, the American Association of Educators, and the Academy of Nutrition and Dietetics. *Diabetes Educ.* 2017;43(1):40-53. doi:10.1177/0145721716689694
 28. Kent D, Haas L, Randal D, et al. Healthy coping: issues and implications in diabetes education and care. *Popul Health Manag.* 2010;13(5):227-233. doi:10.1089/pop.2009.0065
 29. Young-Hyman D, de Groot M, Hill-Briggs F, Gonzalez JS, Hood K, Peyrot M. Psychosocial care for people with diabetes: a position statement of the American Diabetes Association. *Diabetes Care.* 2016;39(12):2126-2140. doi:10.2337/dc16-2053
 30. Fisher EB, Thorpe CT, Devellis BM, Devellis RF. Healthy coping, negative emotions, and diabetes management: a systematic review and appraisal. *Diabetes Educ.* 2007;33(6):1080-1103. doi:10.11770145721707309808
 31. Thorpe CT, Fahey LE, Johnson H, Deshpande M, Thorpe JM, Fisher EB. Facilitating healthy coping in patients with diabetes: a systematic review. *Diabetes Educ.* 2013;39(1):33-52. doi:10.11770145721712464400
 32. Pedersen SS, von Kanel R, Tully PJ, Denollet J. Psychosocial perspectives in cardiovascular disease. *Eur J Prev Cardiol.* 2017;24(3_suppl):108-115. doi:10.11772047487317703827
 33. Shortell SM, Poon BY, Ramsay PP, et al. A multilevel analysis of patient engagement and patient-reported outcomes in primary care practices of accountable care organizations. *J Gen Intern Med.* 2017;32(6):640-647. doi:10.1007/s11606-016-3980-z
 34. Nicolucci A, Kovacs Burns K, Holt RI, et al. Diabetes attitudes, wishes and needs second study (DAWN2): cross-national benchmarking of diabetes-related psychosocial outcomes for people with diabetes. *Diabet Med.* 2013;30(7):767-777. doi:10.1111/dme.12245
 35. Berry E, Lockhart S, Davies M, Lindsay JR, Dempster M. Diabetes distress: understanding the hidden struggles of living with diabetes and exploring intervention strategies. *Postgrad Med J.* 2015;91(1075):278-283. doi:10.1136/postgradmedj-2014-133017
 36. Fisher L, Hessler D, Glasgow RE, et al. REDEEM: a pragmatic trial to reduce diabetes distress. *Diabetes Care.* 2013;36(9):2551-2558. doi:10.2337/dc12-2493
 37. Gonzalvo JD, Hamm J, Eaves S, et al. A practical approach to mental health for diabetes educators. *AADE Pract.* 2019;7(2):29-44. doi:10.11772325160319826929

38. Beckerle CM, Lavin MA. Association of self-efficacy and self-care with glycemic control in diabetes. *Diabetes Spectr*. 2013; 26(3):172-178. doi: 10.2337/diaspect.26.3.172
39. Adam J, Folds L. Depression, self-efficacy, and adherence in patients with type 2 diabetes. *J Nurse Pract*. 2014;10(9):646-652. doi:10.1016/j.nurpra.2014.07.033
40. Fisher L, Hessler DM, Polonsky WH, Mullan J. When is diabetes distress clinically meaningful? Establishing cut points for the Diabetes Distress Scale. *Diabetes Care*. 2012;35(2):259-264. doi:10.2337/dc11-1572
41. Devarajoo C, Chinna K. Depression, distress and self-efficacy: the impact on diabetes self-care practices. *PLoS One*. 2017;12(3): e0175096. doi:10.1371/journal.pone.0175096
42. Karimy M, Koohestani HR, Araban M. The association between attitude, self-efficacy, and social support and adherence to diabetes self-care behavior. *Diabetol Metab Syndr*. 2018;10:86. doi:10.1186/s13098-018-0386-6
43. Indelicato L, Dauriz M, Santi L, et al. Psychological distress, self-efficacy and glycemic control in type 2 diabetes. *Nutr Metab Cardiovasc Dis*. 2017;27(4):300-306. doi.org/10.1016/j.numecd.2017.01.006
44. Aikens JE. Prospective associations between emotional distress and poor outcomes in type 2 diabetes. *Diabetes Care*. 2012;35(12):2472-2478. doi.org/10.2337/dc12-0181
45. Pillay J, Armstrong MJ, Butalia S, et al. Behavioral programs for type 2 diabetes mellitus: a systematic review and network meta-analysis behavioral programs for type 2 diabetes mellitus. *Ann Intern Med*. 2015;163(11):848-860. doi:10.7326/M15-1400
46. Mazaika PK, Weinzimer SA, Mauras N, et al. Variations in brain volume and growth in young children with type 1 diabetes. *Diabetes*. 2016;65(2):476-485. doi:10.2337/db15-1242
47. Cato A, Hershey T. Cognition and type 1 diabetes in children and adolescents. *Diabetes Spectr*. 2016;29(4):197-202. doi:10.2337/ds16-0036
48. Nunley KA, Rosano C, Ryan CM, et al. Clinically relevant cognitive impairment in middle-aged adults with childhood-onset type 1 diabetes. *Diabetes Care*. 2015;38(9):1768-1776. doi:10.2337/dc15-0041
49. Moheet A, Mangia S, Seaquist ER. Impact of diabetes on cognitive function and brain structure. *Ann N Y Acad Sci*. 2015;1353:60-71. doi:10.1111/nyas.12807
50. Munshi MN. Cognitive dysfunction in older adults with diabetes: what a clinician needs to know. *Diabetes Care*. 2017;40(4):461-467. doi:10.2337/dc16-1229
51. Wolf MS, Curtis LM, Wilson EA, et al. Literacy, cognitive function, and health: results of the LitCog study. *J Gen Intern Med*. 2012;27(10):1300-1307. doi:10.1007/s11606-012-2079-4
52. Piatt GA, Rodgers EA, Xue L, Zgibor JC. Integration and utilization of peer leaders for diabetes self-management support: results from project SEED (Support, Education, and Evaluation in Diabetes). *Diabetes Educ*. 2018;44(4):373-382. doi:10.1177/0145721718777855
53. Tang TS, Funnell MM, Sinco B, Spencer MS, Heisler M. Peer-led, empowerment-based approach to self-management efforts in diabetes (PLEASED): a randomized controlled trial in an African American community. *Ann Fam Med*. 2015;13(Suppl 1):S27-S35. doi:10.1370afm.1819
54. Krishnamoorthy Y, Sakthivel M, Sarveswaran G, Eliyas SK. Effectiveness of peer led intervention in improvement of clinical outcomes among diabetes mellitus and hypertension patients—a systematic review and meta-analysis. *Prim Care Diabetes*. 2018;13(2):158-169. doi:10.1016/j.pcd.2018.11.007
55. Domingo JB, Gavero G, Braun KL. Strategies to increase Filipino American participation in cardiovascular health promotion: a systematic review. *Prev Chronic Dis*. 2018;15 (E59):1-11. doi:10.5888/pcd15.170294
56. Heisler M; California HealthCare Foundation. Building peer support programs to manage chronic disease: seven models for success, 2006. Accessed November 2, 2020. <https://www.chcf.org/wp-content/uploads/2017/12/PDF-BuildingPeerSupportPrograms.pdf>
57. Hessler D, Fisher L, Polonsky W, Johnson N. Understanding the areas and correlates of diabetes-related distress in parents of teens with type 1 diabetes. *J Pediatr Psychol*. 2016;41(7):750-758. doi:10.1093/jpepsy/jsw002
58. Polonsky WH, Fisher L, Hessler D, Johnson N. Emotional distress in the partners of type 1 diabetes adults: worries about hypoglycemia and other key concerns. *Diabetes Technol Ther*. 2016;18(5):292-297. doi:10.1089/dia.2015.0451
59. American Diabetes Association. Facilitating behavior change and well-being to improve health outcomes: standards of medical care in diabetes—2020. *Diabetes Care*. 2020;43(Suppl 1):S48-S65. doi:10.2337/dc20-S005
60. Sauter C. Healthy eating. In: Cornell S, Halstenson C, Miller DK, eds. *The Art and Science of Diabetes Care and Education*. 5th ed. Association of Diabetes Care & Education Specialists; 2021:157-182.
61. Academy of Nutrition and Dietetics. Position of the Academy of Nutrition and Dietetics: the role of medical nutrition therapy and registered dietitian nutritionist in the prevention and treatment of prediabetes and type 2 diabetes. *J Acad Nutr Diet*. 2018;118:343-353. doi:10.1016/j.jand.2017.11.021
62. Office of Disease Prevention and Health Promotion. Dietary guidelines 2015-2020. Accessed November 2, 2020. <https://health.gov/dietaryguidelines/2015/guidelines/executive-summary>
63. Hills AP, Byrne NM, Lindstrom R, Hill JO. 'Small changes' to diet and physical activity behaviors for weight management. *Obes Facts*. 2013;6(3):228-238. doi:10.1159/000345030
64. Evert AB, Dennison M, Gardner CD, et al. Nutrition therapy for adults with diabetes or prediabetes: a consensus report. *Diabetes Care*. 2019;42(5):731-754. doi:10.2337/dci19-0014
65. Bowen ME, Cavanaugh KL, Wolff K, et al. The diabetes nutrition education study randomized controlled trial: a comparative effectiveness study of approaches to nutrition in diabetes self-management education. *Patient Educ Couns*. 2016;99(8):1368-1376. doi:10.1016/j.pec.2016.03.017
66. Gibson AA, Hsu MS, Rangan AM, et al. Accuracy of hands v. Household measures as portion size estimation aids. *J Nutr Sci*. 2016;5:e29. doi:10.1017/jns.2016.22
67. Barnard ND, Levin SM, Gloede L, Flores R. Turning the waiting room into a classroom: weekly classes using a vegan or a portion-controlled eating plan improve diabetes control in a randomized translational study. *J Acad Nutr Diet*. 2018;118(6):1072-1079. doi:10.1016/j.jand.2017.11.017
68. Tinker LF, Rosal MC, Young AF, et al. Predictors of dietary change and maintenance in the Women's Health Initiative dietary modification trial. *J Acad Nutr Diet*. 2007;107(7):1155-1166. doi:10.1016/j.jada.2007.04.010
69. Wei M, Gibbons LW, Kampert JB, et al. Low cardiorespiratory fitness and physical inactivity as predictors of mortality in men with type 2 diabetes. *Ann Intern Med*. 2000;132(8):605-611. doi:10.7326/0003-4819-132-8-200004180-00002

70. Powell KE, King AC, Buchner DM, et al. The scientific foundation for the physical activity guidelines for Americans, 2nd edition. *J Phys Act Health*. 2018;16(1):1-11. doi:10.1123/jpah.2018-0618
71. Colberg SR, Sigal RJ, Yardley JE, et al. Physical activity/exercise and diabetes: a position statement of the American Diabetes Association. *Diabetes Care*. 2016;39(11):2065-2079. doi:10.2337/dc16-1728
72. Liu Y, Ye W, Chen Q, et al. Resistance exercise intensity is correlated with attenuation of HbA1c and insulin in patients with type 2 diabetes: a systematic review and meta-analysis. *Int J Environ Res Public Health*. 2019;16(1):140. doi:10.3390/ijerph16010140
73. Wolff-Hughes DL, Fitzhugh EC, Bassett DR, Churilla JR. Total activity counts and bout minutes of moderate-to-vigorous physical activity: relationships with cardiometabolic biomarkers using 2003-2006 NHANES. *J Phys Act Health*. 2015;12(5):694-700. doi:10.1123/jpah.2013-0463
74. Ponsonby A-L, Sun C, Koumoundou OC, et al. Objectively measured physical activity and the subsequent risk of incident dysglycemia: the Australian Diabetes, Obesity and Lifestyle Study (AusDiab). *Diabetes Care*. 2011;34(7):1497-1502. doi:10.2337/dc10-2386
75. Yates T, Davies MJ, Haffner SM, et al. Physical activity as a determinant of fasting and 2-h post-challenge glucose: a prospective cohort analysis of the NAVIGATOR trial. *Diabet Med*. 2015;32(8):1090-1096. doi:10.1111/dme.12762
76. Sallis R, Franklin B, Joy L, et al. Strategies for promoting physical activity in clinical practice. *Prog Cardiovasc Dis*. 2015;57(4):375-386. doi:10.1016/j.pcad.2014.10.003
77. Avery L, Flynn D, Dombrowski SU, et al. Successful behavioural strategies to increase physical activity and improve glucose control in adults with type 2 diabetes. *Diabet Med*. 2015;32(8):1058-1062. doi:10.1111/dme.12738
78. Kärmeniemi M, Lankila T, Ikäheimo T, et al. The built environment as a determinant of physical activity: a systematic review of longitudinal studies and natural experiments. *Ann Behav Med*. 2018;52(3):239-251. doi:10.1093/abm/kax043
79. den Braver NR, Lakerveld J, Rutters F, et al. Built environmental characteristics and diabetes: a systematic review and meta-analysis. *BMC Med*. 2018;16(1):12. doi:10.1186/s12916-017-0997-z
80. Lindsay Smith G, Banting L, Eime R, et al. The association between social support and physical activity in older adults: a systematic review. *Int J Behav Nutr Phys Act*. 2017;14(1):56. doi:10.1186/s12966-017-0509-8
81. Hasan S, Shaw SM, Gelling LH, et al. Exercise modes and their association with hypoglycemia episodes in adults with type 1 diabetes mellitus: a systematic review. *BMJ Open Diabetes Res Care*. 2018;6(1):e000578. doi:10.1136/bmjdr-2018-000578
82. Brazeau AS, Rabasa-Lhoret R, Strychar I, Mircescu H. Barriers to physical activity among patients with type 1 diabetes. *Diabetes Care*. 2008;31(11):2108-2109. doi:10.2337/dc08-0720
83. Kaminsky LA, Dewey D. The association between body mass index and physical activity, and body image, self esteem and social support in adolescents with type 1 diabetes. *Can J Diabetes*. 2014;38(4):244-249. doi:10.1016/j.jcjd.2014.04.005
84. Capoccia K, Odegard PS, Letassy N. Medication adherence with diabetes medication: a systematic review of the literature. *Diabetes Educ*. 2016;42(1):34-71. doi:10.1177/0145721715619038
85. Gallwitz B, Bretzel RG. How do we continue treatment in patients with type 2 diabetes when therapeutic goals are not reached with oral antidiabetes agents and lifestyle? Incretin versus insulin treatment. *Diabetes Care*. 2013;36 (Suppl 2): S180-S189. doi:10.2337/dcS13-2012
86. Polonsky WH, Henry RR. Poor medication adherence in type 2 diabetes: recognizing the scope of the problem and its key contributors. *Patient Prefer Adher*. 2016;10:1299-1307. doi:10.2147PPA.S106821
87. Morello C, Hirsch J. Strategies for addressing the cost of non-adherence in diabetes. *Am J Manag Care*. 2017;23:S247-S252.
88. Cramer JA, Roy A, Burrell A, et al. Medication compliance and persistence: terminology and definitions. *Value Health*. 2008; 11(1):44-47. doi:10.1111/j.1524-4733.2007.00213.x
89. World Health Organization. Adherence to long-term therapies: evidence for action, 2003. Accessed November 2, 2020. <https://apps.who.int/iris/bitstream/handle/10665/42682/9241545992.pdf;jsessionid=F16092C144CF5EAA49CB92A1092C707A?sequence=1>
90. American Diabetes Association. Cardiovascular disease and risk management: standards of medical care in diabetes—2020. *Diabetes Care*. 2020;43(Suppl 1):S11-S134. doi:10.2337/dc20-s010
91. DeFronzo RA. Banting Lecture. From the triumvirate to the ominous octet: a new paradigm for the treatment of type 2 diabetes mellitus. *Diabetes*. 2009;58:773-795. doi:10.2337/db09-9028
92. Cavaiaola TS, Pettus JH. Management of type 2 diabetes: selecting amongst available pharmacological agents. In: Feingold KR, Anawalt B, Boyce A, et al., eds. *Endotext* [Internet]. MDText.com, Inc.; 2000.
93. Stadler LKJ, Farooqi IS. A new drug target for type 2 diabetes. *Cell*. 2017;170(1):12-14. doi:10.1016/j.cell.2017.06.024
94. Miller BR, Nguyen H, Hu CJ, Lin C, Nguyen QT. New and emerging drugs and targets for type 2 diabetes: reviewing the evidence. *Am Health Drug Benefits*. 2014;7(8):452-463.
95. Lipska KJ, Krumholz HM. Is hemoglobin A1c the right outcome for studies of diabetes? *JAMA*. 2017;317:1017-1018. doi:10.1001/jama.2017.0029
96. Richard MB. Glycemic variability and diabetes complications: does it matter? Simply put, there are better glycemic markers! *Diabetes Care*. 2015;38(8):1615-1621. doi:10.2337/dc15-0099
97. Hirsch IB, Brownlee M. Beyond hemoglobin A1c—need for additional markers of risk for diabetic microvascular complications. *JAMA*. 2010;303:2291-2292. doi:10.1001/jama.2010.785
98. Ismail-Beigi F, Moghissi E, Tiktin M, Hirsch IB, Inzucchi SE, Genuth S. Individualizing glycemic targets in type 2 diabetes mellitus: implications of recent clinical trials. *Ann Intern Med*. 2011;154:554-549. doi:10.7326/0003-4819-154-8-201104190-00007
99. Inzucchi SE, Bergenstal RM, Buse JB, et al. Management of hyperglycemia in type 2 diabetes: a patient-centered approach: position statement of the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetes Care*. 2012;35:1364-1379. doi:10.2337/dc12-0413
100. Hunt CW. Technology and diabetes self-management: an integrative review. *World J Diabetes*. 2015;6(2):225-233. doi:10.4239/wjd.v6.i2.225
101. Fitipaldi H, McCarthy MI, Florez JC, Franks PW. A global overview of precision medicine in type 2 diabetes. *Diabetes*. 2018;67:1911-1922. doi:10.2337/dbi17-0045

102. Azzi M, Constantino M, Pont L, McGill M, Twigg S, Krass I. Medication safety: an audit of medication discrepancies in transferring type 2 diabetes mellitus (T2DM) patients from Australian primary care to tertiary ambulatory care. *Int J Qual Health Care*. 2014;26(4):397-403. doi:10.1093/intqhc/mzu051
103. Turchin A, Sosina O, Zhang H, et al. Ambulatory medication reconciliation and frequency of hospitalizations and emergency department visits in patients with diabetes. *Diabetes Care*. 2018;41(8):1639-1645. doi:10.2337/dc17-1260
104. Sofaer S. Navigating poorly charted territory: patient dilemmas in health care "nonsystems." *Med Care Res Rev*. 2009;66(Suppl 1):75S-93S. doi:10.1177/1077558708327945
105. Polonsky WH, Skinner TC. Perceived treatment efficacy: an overlooked opportunity in diabetes care. *Clin Diabetes*. 2010;28(2):89-92. doi:10.2337/diaclin.28.2.89
106. Hajos TR, Polonsky WH, Pouwer F, Gonder-Frederick L, Snoek FJ. Toward defining a cutoff score for elevated fear of hypoglycemia on the hypoglycemia fear survey worry subscale in patients with type 2 diabetes. *Diabetes Care*. 2014;37(1):102-108. doi:10.2337/dc13-0971
107. Gonzalez JS, Schenider HE, Wexler DJ, et al. Validity of medication adherence self-reports in adults with type 2 diabetes. *Diabetes Care*. 2013;36:831-837. doi:10.2337/dc12-0410
108. Zullig LL, Bosworth H. Engaging patients to optimize medication adherence. *NEJM Catal*. May 14, 2017. Accessed November 2, 2020. <https://catalyst.nejm.org/optimize-patients-medication-adherence/>
109. American Association of Diabetes Educators. The role of diabetes educator in diabetes formulary and medical device decisions. *Diabetes Educ*. 2019;45(1):50-53. doi:10.1177/0145721718820942
110. American Association of Diabetes Educators. The role of diabetes educator in inpatient diabetes management. *Diabetes Educ*. 2017;43(1):28-33. doi:10.1177/0145721718754817
111. Piwek L, Ellis DA, Andrews S, Joinson A. The rise of consumer health wearables: promises and barriers. *PLoS Med*. 2016;13(2):e1001953. doi:10.1371/journal.pmed.1001953
112. Blackwell M, Wheeler BJ. Clinical review: the misreporting of logbook, download, and verbal self-measured blood glucose in adults and children with type 1 diabetes. *Acta Diabetol*. 2017;54(1):1-8. doi:10.1007/s00592-016-0907-4
113. Danne T, Nimri R, Battelino T, et al. International consensus on use of continuous glucose monitoring. *Diabetes Care*. 2017;40(12):1631-1640. doi:10.2337/dc17-1600
114. Rao GHR. Integrative approach to the management of cardiometabolic diseases. *J Cardiol Cardiovasc Sci*. 2018;2(3):37-42.
115. Dayer L, Heldenbrand S, Anderson P, Gubbins PO, Martin BC. Smartphone medication adherence apps: potential benefits to patients and providers. *J Am Pharm Assoc*. 2013;53(2):172-181. doi:10.1331/JAPhA.2013.13121
116. Polonsky WH, Fisher L, Schikman CH, et al. Structured self-monitoring of blood glucose significantly reduces A1C levels in poorly controlled, noninsulin-treated type 2 diabetes results from the structured testing program study. *Diabetes Care*. 2011;34(2):262-267. doi:10.2337/dc10-1732
117. Polonsky WH, Hessler D, Ruedy KJ, et al. The impact of continuous glucose monitoring on markers of quality of life in adults with type 1 diabetes: further findings from the Diamond randomized clinical trial. *Diabetes Care*. 2017;40(6):736-741. doi:10.2337/dc17-0133
118. McManus RJ, Mant J, Haque MS, et al. Effect of self-monitoring and medication self-titration on systolic blood pressure in hypertensive patients at high risk of cardiovascular disease: the TASMIN-SR randomized clinical trial. *JAMA*. 2014;312(8):799-808. doi:10.1001/jama.2014.10057
119. The Office of the National Coordinator for Health Information Technology. Patient-generated health data. Accessed November 2, 2020. <https://www.healthit.gov/topic/scientific-initiatives/patient-generated-health-data>
120. DeCoste KC, Miller DK. Reducing risks. In: Cornell S, Halstenon C, Miller DK, eds. *The Art and Science of Diabetes Care and Education*. 5th ed. Association of Diabetes Care & Education Specialists; 2021:285-306.
121. American Diabetes Association. Introduction: standards of medical care in diabetes—2020. *Diabetes Care*. 2020;43(Suppl 1):S1-S2.
122. Gopalan A, Lorincz IS, Wirtalla C, Marcus SC, Long JA. Awareness of prediabetes and engagement in diabetes risk-reducing behaviors. *Am J Prev Med*. 2015;49(4):512-519. doi:10.1016/j.amepre.2015.03.007
123. DiBenedetto JC, Blum NM, O'Brian CA, Kolb LE, Lipman RD. Achievement of weight loss and other requirements of the diabetes prevention and recognition program: a National Diabetes Prevention Program network based on nationally certified diabetes self-management education programs. *Diabetes Educ*. 2016;42(6):678-685. doi:10.1177/0145721716668415
124. Knowler WC, Fowler SE, Hamman RF, et al. 10-year follow-up of diabetes incidence and weight loss in the Diabetes Prevention Program Outcomes Study. *Lancet*. 2009;374(9702):1677-1686. doi:10.1016/S0140-6736(09)61457-4
125. Mudaliar U, Zabetian A, Goodman M, et al. Cardiometabolic risk factor changes observed in diabetes prevention programs in US settings: a systematic review and meta-analysis. *PLoS Med*. 2016;13(7):e1002095. doi:10.1371/journal.pmed.1002095
126. Ely E, Gruss S, Luman E, et al. A national effort to prevent type 2 diabetes: participant-level evaluation of CDC's National Diabetes Prevention Program. *Diabetes Care*. 2017;40(10):1331-1341. doi:10.2337/dc16-2099
127. Chvala CA, Sherr D, Lipman RD. Diabetes self-management education for adults with type 2 diabetes mellitus: a systematic review of the effect on glycemic control. *Patient Educ Couns*. 2016;99(6):926-943. doi:10.1016/j.pec.2015.11.003
128. Nathan DM, Genuth S, Lachin J, et al. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med*. 1993;329(14):977-986. doi:10.1056/NEJM199309303291401
129. UK Prospective Diabetes Study (UKPDS) Group. Effect of intensive blood-glucose control with metformin on complications in overweight patients with type 2 diabetes (UKPDS 34). *Lancet*. 1998;352(9131):854-865. doi:10.1016/S0140-6736(98)07037-8
130. UK Prospective Diabetes Study (UKPDS) Group. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet*. 1998;352(9131):837-853. doi:10.1016/S0140-6736(98)07019-6
131. Steinsbekk A, Rygg LO, Lisulo M, Rise MB, Fretheim A. Group based diabetes self-management education compared to routine treatment for people with type 2 diabetes mellitus. A

- systematic review with meta-analysis. *BMC Health Serv Res.* 2012;12:213. doi:10.1186/1472-6963-12-213
132. Duncan I, Ahmed T, Li QE, et al. Assessing the value of the diabetes educator. *Diabetes Educ.* 2011;37(5):638-657. doi:10.11770145721711416256
133. Duncan I, Birkmeyer C, Coughlin S, Li QE, Sherr D, Boren S. Assessing the value of diabetes education. *Diabetes Educ.* 2009;35(5):752-760. doi:10.11770145721709343609
134. Johnson KA, Chen S, Cheng IN, et al. The impact of clinical pharmacy services integrated into medical homes on diabetes-related clinical outcomes. *Ann Pharmacother.* 2010;44(12):1877-1886. doi:10.1345aph.1P380
135. Lee SWH, Ng KY, Chin WK. The impact of sleep amount and sleep quality on glycemic control in type 2 diabetes: a systematic review and meta-analysis. *Sleep Med Rev.* 2017;31:91-101. doi:10.1016/j.smrv.2016.02.001
136. American Diabetes Association. Improving care and promoting health in populations: standards of medical care in diabetes—2020. *Diabetes Care.* 2020;43(Suppl 1):S7-S13. doi:10.2337/dc20-S001
137. American Diabetes Association. Comprehensive medical evaluation and assessment of comorbidities: standards of medical care in diabetes—2020. *Diabetes Care.* 2020;43(Suppl 1):S37-S47. doi:10.2337/dc20-s004
138. American Diabetes Association. Microvascular complications and foot care: standards of medical care in diabetes—2020. *Diabetes Care.* 2020;43(Suppl 1):S135-S151. doi:10.2337/dc20-s011
139. King DK, Glasgow RE, Toobert DJ, et al. Self-efficacy, problem solving, and social-environmental support are associated with diabetes self-management behaviors. *Diabetes Care.* 2010;33(4):751-753. doi:10.2337/dc09-1746
140. Mensing C. Problem solving. In: Cornell S, Halstenson C, Miller DK, eds. *The Art and Science of Diabetes Care and Education*. 5th ed. Association of Diabetes Care & Education Specialists; 2021:307-326.
141. Hill-Briggs F, Gemmell L. Problem solving in diabetes self-management and control: a systematic review of the literature. *Diabetes Educ.* 2007;33(6):1032-1050. doi:10.1177/0145721707308412
142. Fitzpatrick SL, Schumann KP, Hill-Briggs F. Problem solving interventions for diabetes self-management and control: a systematic review of the literature. *Diabetes Res Clin Pract.* 2013;100(2):145-161. doi.org/10.1016/j.diabres.2012.12.016
143. Hill-Briggs F, Lazo M, Peyrot M, et al. Effect of problem-solving-based diabetes self-management training on diabetes control in a low income patient sample. *J Gen Intern Med.* 2011;26(9):972-978. doi.org/10.1007/s11606-011-1689-6
144. Fitzpatrick SL, Golden SH, Stewart K, et al. Effect of DECIDE (Decision-making Education for Choices In Diabetes Everyday) program delivery modalities on clinical and behavioral outcomes in urban African Americans with type 2 diabetes: a randomized trial. *Diabetes Care.* 2016;39(12):2149-2157. doi.org/10.2337/dc16-0941
145. Lilly CL, Bryant LL, Leary JM, et al. Evaluation of the effectiveness of a problem-solving intervention addressing barriers to cardiovascular disease prevention behaviors in 3 underserved populations: Colorado, North Carolina, West Virginia, 2009. *Prev Chronic Dis.* 2014;11:E32. doi:10.5888/pcd11.130249
146. Bodenheimer T, Lorig K, Holman H, Grumbach K. Patient self-management of chronic disease in primary care. *JAMA.* 2002;288(19):2469-2475. doi:10.1001/jama.288.19.2469
147. Centers for Disease Control and Prevention. Talking points about health literacy, 2016. Accessed November 2, 2020. <https://www.cdc.gov/healthliteracy/shareinteract/TellOthers.html>
148. Serrano V, Rodriguez-Gutierrez R, Hargraves I, Gionfriddo MR, Tamhane S, Montori VM. Shared decision-making in the care of individuals with diabetes. *Diabet Med.* 2016;33(6):742-751. doi:10.1111/dme.13143
149. Miller CK, Bauman J. Goal setting: an integral component of effective diabetes care. *Curr Diab Rep.* 2014;14(8):509. doi:10.1007/s11892-014-0509-x
150. Fredrix M, McSharry J, Flannery C, Dinneen S, Byrne M. Goal-setting in diabetes self-management: a systematic review and meta-analysis examining content and effectiveness of goal-setting interventions. *Psychol Health.* 2018;33(8):955-977. doi:10.1080/08870446.2018.1432760
151. O'Donnell M, Carey ME, Horne R, et al. Assessing the effectiveness of a goal-setting session as part of a structured group self-management education programme for people with type 2 diabetes. *Patient Educ Couns.* 2018;101(12):2125-2133. doi.org/10.1016/j.pec.2018.07.009
152. Center for Disease Control and Prevention. Social determinants of health: know what affects health. Accessed November 2, 2020. <https://www.cdc.gov/socialdeterminants/faqs/index.htm>
153. Fisher EB, Brownson CA, O'Toole ML, Shetty G, Anwuri VV, Glasgow RE. Ecological approaches to self-management: the case of diabetes. *Am J Public Health.* 2005;95(9):1523-1535. doi:10.2105/AJPH.2005.066084
154. Nester J. The importance of interprofessional practice and education in the era of accountable care. *NC Med J.* 2016;77(2):128-132. doi:10.18043/ncm.77.2.128
155. Schottenfeld L, Petersen D, Peikes D, et al; Agency for Healthcare Research and Quality. Creating patient-centered team-based primary care. AHRQ Publication No. 16-0002-EF. 2016. Accessed November 27, 2020. <https://pcmh.ahrq.gov/page/creating-patient-centered-team-based-primary-care>
156. Okun S, Schoenbaum S, Andrews D, et al. Patients and health care teams forging effective partnerships. December 17, 2014. <https://nam.edu/perspectives-2014-patients-and-health-care-teams-forging-effective-partnerships/> doi:10.31478/201412f
157. Bodenheimer T, Sinsky C. From triple to quadruple aim: care of the patient requires care of the provider. *Ann Fam Med.* 2014;12(6):573-576. doi:10.1370/afm.1713

For reprints and permission queries, please visit SAGE's Web site at <https://us.sagepub.com/en-us/nam/journals-permissions>.